

Hasani Swindell, M.D.

Sports Medicine
Shoulder, Elbow & Knee Arthroscopy
Shoulder Replacement Surgery
T: 718.246.8700
F: 718.246.8701

POSTOPERATIVE INSTRUCTIONS

Patellar Tendon Repair

Recovery after knee surgery entails healing, controlling swelling and discomfort, return of range of motion of the knee joint, regaining strength in the muscles around the knee joint and a gradual return to activities. The following instructions are intended as a guide to help you achieve these goals and recovery as quickly as possible after your knee surgery.

DIET

- Your first few meals after surgery should include light, easily digestible foods and plenty of liquids, since some people experience nausea as a temporary reaction to anesthesia
- Progress to your normal diet if you are not nauseated

WOUND CARE

- Maintain your operative dressing, loosen bandage if swelling of the foot and ankle occurs.
- It is normal for the knee to bleed and swell following surgery. If blood soaks onto the ACE bandage, do not become alarmed, reinforce with additional dressing.
- To avoid infection, keep surgical incisions clean and dry for the first 7 days following surgery – you may shower by placing a large plastic bag over your brace beginning the day after surgery. NO immersion of the operative leg (ie: bath or pool).
- You may remove the outer surgical dressings on the 7th post-operative day
 - Leave the clear, waterproof dressing intact until your follow-up appointment
- Please do not place any ointments lotions or creams directly over the incisions.
- Once the sutures are removed at least 7–10 days post operatively you can begin to get the incision wet in the shower (water and soap lightly run over the incision and pat dry). NO immersion in a bath until given approval by our office.

MEDICATIONS

- Local anesthetics are injected into the wound and knee joint at the time of surgery. This will wear off within 8–12 hours and it is not uncommon for patients to encounter more pain on the first or second day after surgery when swelling peaks.
- Most patients will require some narcotic pain medication for a short period of time – this can be taken as per directions on the bottle.
 - **Narcotic prescriptions may be refilled up to 2 times**

Hasani Swindell, M.D.

Sports Medicine
Shoulder, Elbow & Knee Arthroscopy
Shoulder Replacement Surgery

T: 718.246.8700

F: 718.246.8701

- Common side effects of the pain medication are nausea, drowsiness, and constipation. To decrease the side effects take the medication with food. If constipation occurs, consider taking an over the counter laxative.
- If you are having problems with nausea and vomiting, contact the office to possibly have your medications changed.
- Do not drive a car or operate machinery while taking the narcotic medication.
- Please avoid alcohol use while taking narcotic pain medication
- If you are having pain that is not being controlled by the pain medication prescribed, you may take an over the counter anti-inflammatory medication such as ibuprofen or naproxen in between doses of pain medication. This will help to decrease pain and decrease the amount of narcotic medication required. Please take as directed on the bottle.
- For 3 weeks following surgery take one aspirin 325mg tablet twice daily to lower the risk of developing a blood clot after surgery. Please contact the office should severe calf pain occur or significant swelling of the calf or ankle occur

ACTIVITY

- Elevate the operative leg to chest level whenever possible to decrease swelling.
- Do not place pillows under knees (i. e. do not maintain knee in a flexed or bent position), but rather place pillows under the foot /ankle.
- Use crutches to assist with walking – WBAT with the knee locked in extension. You must weightbear with the brace on locked in full extension after surgery. Do not walk without brace on.
- Do not engage in activities which increase knee pain /swelling (prolonged periods of standing or walking) for the first 7-10 days following surgery.
- Avoid long periods of sitting (without leg elevated) or long distance traveling for 2 weeks.
- NO driving until instructed otherwise by physician.
- May return to sedentary work ONLY or school 3-4 days after surgery, if pain is tolerable.
- Goals for motion are:
 - 0-30 degrees – Weeks 2-4
 - 0-60 degrees – Weeks 4-6
 - 0-90 degrees – Weeks 6-8

BRACE

- Your brace should be worn fully extended (straight) at all times (day and night – except for exercises) until otherwise instructed after the first post-operative visit

Hasani Swindell, M.D.

Sports Medicine
Shoulder, Elbow & Knee Arthroscopy
Shoulder Replacement Surgery

T: 718.246.8700

F: 718.246.8701

ICE THERAPY

- Icing is very important in the initial post-operative period and should begin immediately after surgery.
- Use icing machine continuously or ice packs (if machine not prescribed) for 30–45 minutes every 2 hours daily until your first post-operative visit – remember to keep leg elevated to level of chest while icing. Care should be taken with icing to avoid frostbite to the skin. Keep some barrier (clothing, towel, dressing) between the ice and your bare skin
- You do not need to wake up in the middle of the night to change over the ice machine or icepacks unless you are uncomfortable.

EXERCISE

- Begin exercises 24 hours after surgery (straight leg raises, quad sets, heel slides, and ankle pumps) unless otherwise instructed. (See pictures of exercises below for reference)
- Discomfort and knee stiffness is normal for a few days following surgery. It is safe to bend your knee in a non-weightbearing position when performing exercises unless otherwise instructed. Avoid flexing past 90 degrees per goals listed above
- Complete exercises 3–4 times daily until your first post-operative visit – your motion goals are to have complete extension (straightening) and 90 degrees of flexion (bending) at your 6 to 8 week appointment
- Perform ankle pumps continuously throughout the day to reduce the risk of developing a blood clot in your calf.
- Formal physical therapy (PT) typically begins 2 weeks after surgery. A prescription and protocol will be provided at your first post-op visit.

EMERGENCIES

- Contact Dr. Swindell's team at 718-246-8700 if any of the following are present:
 - Painful swelling or numbness (note that some swelling and numbness is normal).
 - Unrelenting pain.
 - Fever (over 101° – it is normal to have a low grade fever for the first day or two following surgery) or chills.
 - Redness around incisions.
 - Color change in foot or ankle.
 - Continuous drainage or bleeding from incision (a small amount of drainage is expected).
 - Difficulty breathing
 - Excessive nausea /vomiting.

Hasani Swindell, M.D.

Sports Medicine

Shoulder, Elbow & Knee Arthroscopy

Shoulder Replacement Surgery

T: 718.246.8700

F: 718.246.8701

- Calf pain
- If you have an emergency after office hours or on the weekend, contact the office at 718-246-8700 and you will be connected to our answering service.
 - If you are enrolled in Epic Connect, you can also reach the office through the patient portal
- If you have an emergency that requires immediate attention proceed to the nearest emergency room.

FOLLOW-UP CARE /QUESTIONS

- If you do not already have a post-operative appointment scheduled, please contact our scheduler at 718-246-8700 schedule.
- Typically the first post-operative appointment following surgery is 10-14 days following surgery.
- At your first post operative appointment, you will have a wound check, go over physical therapy protocols and we will answer any questions you may have about the procedure.